



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Plant contributions to our understanding of sex chromosome evolution

Citation for published version:

Charlesworth, D 2015, 'Plant contributions to our understanding of sex chromosome evolution', *New Phytologist*, vol. 208, no. 1, pp. 52–65. <https://doi.org/10.1111/nph.13497>

Digital Object Identifier (DOI):

[10.1111/nph.13497](https://doi.org/10.1111/nph.13497)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

New Phytologist

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





New Phytologist

**Plant contributions to our understanding of sex
chromosome evolution**

Journal:	<i>New Phytologist</i>
Manuscript ID:	NPH-TR-2015-19155.R2
Manuscript Type:	TR - Commissioned Material - Tansley Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Charlesworth, Deborah; University of Edinburgh, School of Biological Sciences;
Key Words:	Sex chromosomes, recombination suppression, partial sex linkag, sex-determination, genetic degeneration

SCHOLARONE™
Manuscripts

Review

1 Tansley review for *New Phytologist*

2

3

4 Title: **Plant contributions to our understanding of sex chromosome evolution**

5 Deborah Charlesworth

6

7 Address for correspondence:

8 Institute of Evolutionary Biology, University of Edinburgh,

9 Ashworth Lab. King's Buildings, W. Mains Rd., Edinburgh EH9 3FL, UK

10 Phone: 0131 650 5751

11 Email: Deborah.Charlesworth@ed.ac.uk

12

13

14 Key words: Sex chromosomes, recombination suppression, partial sex linkage, sex
15 determination, genetic degeneration

16

17

18

19

20

21

22

23

24 **Summary**

25 A minority of angiosperms have male and female flowers separated in distinct
26 individuals (dioecy), and most dioecious plants do not have cytologically different
27 (heteromorphic) sex chromosomes. Plants nevertheless have several advantages
28 for studying sex chromosome evolution, as genetic sex determination has evolved
29 repeatedly and is often absent in close relatives. I review sex-determining regions
30 in non-model plant species, which may help us understand when and how (and,
31 potentially, test hypotheses about why) recombination suppression evolves within
32 young sex chromosomes. I emphasise high throughput sequencing approaches that
33 are increasingly being applied to plants to test for non-recombining regions. These
34 data are particularly illuminating when combined with sequence data that allow
35 phylogenetic analyses, and estimates of when these regions evolved. Together
36 with comparative genetic mapping, this has revealed that sex-determining loci and
37 sex-linked regions evolved independently in many plant lineages, sometimes in
38 closely related dioecious species, and often within the past few million years. In
39 reviewing recent progress, I suggest areas for future work, such as using
40 phylogenies to allow informed choice of outgroup species suitable for inferring
41 the directions of changes, including testing whether Y chromosome-like regions
42 are undergoing genetic degeneration, a predicted consequence of losing
43 recombination.

44

45 **Introduction: Advantages of plants for studying sex chromosome evolution**

46 Most flowering plants have hermaphroditic flowers, and only a minority have
47 separate male and female flowers (monoecy or dioecy). Among dioecious plants, with
48 male and female flowers separated in distinct individuals, some species have
49 environmental, not genetic, control of sex determination (Policansky, 1981;
50 Zimmerman, 1991; Pannell, 1997), and those with genetic sex determination often do
51 not have cytologically differentiated sex chromosomes (Westergaard, 1958; Ming *et al.*,
52 2011; Renner, 2014). In contrast, separate sexes and heteromorphic sex
53 chromosomes are common in many familiar animal groups (Bachtrog, 2012).
54 Nevertheless, plants have several advantages for research on sex chromosomes,
55 because genetic sex determination has evolved repeatedly among angiosperms, and
56 independently in different families (Charlesworth, 1985; Ming *et al.*, 2011; Renner,
57 2014). Compared with the best-studied animal systems (Bellott *et al.*, 2014; Cortez *et al.*,
58 2014; Zhou *et al.*, 2014), many flowering plant sex chromosomes probably
59 evolved very recently (Marais *et al.*, 2011; Renner, 2014), yet, as will be illustrated
60 below, similarities with animal systems are striking. Table 1 summarises the main
61 advantages of using dioecious plants to study sex chromosome evolution, and test
62 hypotheses about sex chromosome evolution derived from theoretical modelling.

63
64 Table 1 about here

65
66 There is now too much published work on plant sex chromosomes and their
67 evolution to include in a single article. I therefore focus on recombination
68 suppression, the defining characteristic of sex chromosomes, which leads to the
69 evolution of sex chromosomes' other unusual characteristics, genetic degeneration
70 and accumulation of repetitive sequences on the sex chromosomes, which I mention
71 only briefly. I review progress that has come through genetic and molecular
72 evolutionary studies, illustrating how this has involved combining approaches,
73 including DNA sequencing, and resequencing of multiple individuals of the same sex
74 and species for genetic and population genetic tests of sex-linkage, together with
75 sequencing for phylogenetic studies, and to place events in well-established time
76 frames.

77

78

79 **Sex chromosomes and estimating their ages from sequence divergence**

80 I define plant sex chromosomes as genome regions of these species that carry
 81 the “SEX” locus that controls individuals' sexes, and that do not recombine. Rather
 82 than using the term sex chromosomes, I shall often use “fully sex-linked regions” (and
 83 full sex linkage), because of the diversity among plants with genetically controlled
 84 dioecy — some have extensive non-pairing regions that show heteromorphism
 85 between the sexes, like many animal sex chromosomes, but many have no detectable
 86 cytological differences (recently reviewed by Renner, 2014). *Silene latifolia* is an
 87 example of sex chromosome heteromorphism. It has an XY system, and males are the
 88 heterozygous sex, and, as in mammals (Bellott *et al.*, 2014; Cortez *et al.*, 2014). The
 89 Y is largely non-recombining, with XY pairing only in a small pseudo-autosomal
 90 region (PAR) region at one tip (Westergaard, 1958; Filatov *et al.*, 2008); mapping of
 91 genic markers suggests a single PAR (Bergero *et al.*, 2013), though an AFLP map
 92 suggests two (Scotti & Delph, 2006). However, unlike many animal Y chromosomes,
 93 the fully sex-linked region still carries hundreds of genes (Bergero & Charlesworth,
 94 2011; Chibalina & Filatov, 2011; Muyle *et al.*, 2012). In contrast, the fully Y-linked
 95 region in papaya is only about 10% of chromosome 1 (Liu *et al.*, 2004; Wang *et al.*,
 96 2012). Some diploid plants have ZW systems, in which females are ZW heterozygotes
 97 and males are ZZ homozygotes (Westergaard, 1958), as in birds (Zhou *et al.*, 2014)
 98 and Lepidoptera (Suetsugu *et al.*, 2013); these include *Fragaria* (strawberry) species
 99 (Spigler *et al.*, 2008; Goldberg *et al.*, 2010) and *Salix* (Alstrom-Rapaport *et al.*, 1998).
 100 Other systems, including those in haploid plants, will be described below.

101 The time when recombination stopped can be estimated using DNA sequence
 102 divergence between genes present on the Y as well as the X, together with a
 103 “molecular clock” for synonymous or silent site divergence per year. Higher X-Y
 104 divergence values correspond to greater times since recombination suppression. In
 105 both humans (Lahn & Page, 1999) and the plant *S. latifolia* (Bergero *et al.*, 2007),
 106 divergence increases with the distance from the PAR (in X chromosome genetic or
 107 physical maps; (as these Y chromosomes are extensively rearranged, distances on the
 108 Y are not informative Skaletsky *et al.*, 2003; Bergero *et al.*, 2008). Therefore,
 109 suppressed recombination must have spread from an early non-recombining region,
 110 the oldest “evolutionary stratum” (Lahn & Page, 1999), towards younger “strata”

111 closer to the current PAR. X-Y divergence in the older *S. latifolia* stratum is similar to
 112 that in the youngest of the five strata in humans (Skaletsky *et al.*, 2003), and the
 113 *Silene* XY pair probably evolved about 5-10 MYA (Nicolas *et al.*, 2005).

114 [Figure 1 about here]

115 A sex chromosome system may be older than its oldest stratum, because
 116 recombination suppression in a sex-determining region usually takes time to evolve
 117 (see below). On the other hand, recombination suppression may pre-date evolution of
 118 separate sexes. In several well-studied plants, results from combining genetic and
 119 physical mapping reveal large genome regions with infrequent crossing over
 120 surrounding the centromeres, with crossovers restricted to the ends of chromosomes,
 121 for example in maize (Rodgers-Melnicka *et al.*, 2015). These regions may include
 122 substantial proportions of genes; in barley, for example, about 20% of genes are
 123 estimated to be located in such regions (Baker *et al.*, 2014). If sex-determining loci
 124 evolve in such a region, the oldest stratum will be contemporaneous with the sex-
 125 determination system (Figure 1).

126 In what follows, I stress the importance of estimating ages of non-recombining
 127 regions for understanding several important aspects of sex chromosome evolution.
 128 Young sex chromosome systems are well suited for studying the early stages of
 129 evolution of recombination suppression and the evolution of these characteristics; in
 130 older animal systems, these processes can only be studied over a coarse time scale
 131 that cannot reveal much detail. Young evolutionary strata in plant sex chromosomes
 132 are also of interest for studying the time course of genetic degeneration, including
 133 gene losses from Y chromosomes.

134

135 **Which plants have sex chromosomes?**

136 Genetic maps can detect the presence of sex-linked regions in dioecious species.
 137 In papaya, for example, a large set of AFLP molecular variants was first mapped in a
 138 full-sib family (Liu *et al.*, 2004). The completely sex-linked region is small, making
 139 BAC sequencing of the region possible, which showed that the X-linked region
 140 includes only 3.5 Mb (it is flanked by much larger PARs), and carries about 50 genes
 141 with apparently functional Y-linked copies (Wang *et al.*, 2012). Assembly of the
 142 physical map of the homologous Y-like region suggests that part of the sex-linked

143 region is probably in a pericentromeric region (Yu *et al.*, 2007; Zhang *et al.*, 2008).
144 Silent site divergence values for XY pairs suggests that inversions occurred,
145 suppressing recombination, about 7 and 1.7 million years ago, implying that a new
146 recombination-suppressed region has formed since the older stratum evolved (Figure
147 1).

148 Ideally, sex-linkage in a family should be confirmed by showing that Y-linked
149 variants are found only in males in a wider sample of genotypes (from natural
150 population samples, or from multiple cultivars of crop species) to exclude partially
151 sex-linked genes in the PAR that did not yield recombinants in the particular cross
152 studied. This is unnecessary for papaya, because, although this is clearly a young
153 system, the sequence divergence across part of the X-Y region is around 7% for silent
154 sites, much higher than between alleles in recombining regions of the genome,
155 including the collinear regions adjoining the fully sex-linked sequences (Wang *et al.*,
156 2012). This strongly suggests complete sex-linkage.

157 Now that large numbers of genetic markers can be developed in non-model
158 organisms, using high-throughput approaches, it will be possible to discover how
159 many other plant cases like papaya exist, without major cytologically detectable sex
160 chromosome heteromorphism, but with fully sex-linked regions carrying multiple
161 genes, and assess how many plants evolved dioecy so recently that their sex-
162 determining loci have not yet evolved non-recombining sex chromosome-like regions.
163 I next outline other approaches that have demonstrated that sex-linked regions have
164 evolved in dioecious plants.

165 When divergence data are not available, genetic mapping in related species can
166 help test whether recombination suppression has evolved in dioecious plants. If the
167 SEX locus of a dioecious species is in a genome region of suppressed recombination,
168 but the homologous region of a non-dioecious relative recombines, this would suggest
169 that recombination suppression evolved following the evolution of genetically
170 controlled dioecy, rather than being the ancestral state (unless the SEX locus is in a
171 pericentromeric region whose extent or location has changed between the species). A
172 dioecious close relative of papaya, *Vasconcellea parviflora*, has been shown to have a
173 homologous SEX locus, based on cytogenetic detection of heterochromatin in the
174 centromere-proximal regions of the homologous chromosomes of the two species.
175 This result also shows that the papaya and *V. parviflora* sex chromosomes are not
176 truly homomorphic (Iovene *et al.*, 2015) — their heteromorphism is minor, but

detectable with refined modern cytological methods, consistent with the sequencing results showing that the papaya Y--linked region is larger than the X region (Wang *et al.*, 2012).

These examples illustrate the value of genetic mapping within families. However, studies of wider population samples can also be used to discover sex-linked regions and establish which is the heterozygous sex. In an XY system, all males should be heterozygotes for fully sex-linked alleles, and population surveys can reveal such male-specific variants (which can be confirmed by data on segregation patterns within families). This approach has established that the date palm (*Phoenix dactylifera*) has at least several fully sex-linked genes (Cherif *et al.*, 2012). Neither the physical size of the SEX region, nor the age of this system has yet been estimated. It is a heteromorphic XY system (Siljak-Yakovlev *et al.*, 1996) that may be ancient, as many other species in the palm family are also dioecious (Renner, 2014).

190

191 **Haploid plants**

Many haploid plants have sex chromosomes, as determined either from morphology differences in the karyotypes of male and female gametophytes (Bull, 1983; Ming *et al.*, 2011), or from the existence of one or more sex-linked genetic markers (Immmler & Otto, 2015). The male- and female-determining chromosomes of haploids are now often called V and U, respectively (Bachtrog *et al.*, 2011), to emphasize that the SEX region is never homozygous and can therefore never recombine; until physical maps are produced, it is not possible to relate sequence divergence to the genetic map position in the sex-linked region. The older literature, including studies in *Marchantia polymorpha*, called them X/Y systems (Okada *et al.*, 2001; Yamato *et al.*, 2007). *M. polymorpha* has highly heteromorphic sex chromosomes, and divergence between alleles of the few sex-linked gene pairs studied is extremely high, indicating an ancient system.

In contrast, genetic mapping in the moss *C. purpureus* found that, as in papaya, only markers in the middle of the linkage group with the SEX locus show full sex-linkage (McDaniel *et al.*, 2007). Sequence data for all site types in coding plus (predominantly) non-coding regions of U- and V- linked allele sequences suggests that evolutionary strata may exist in *C. purpureus*. Divergence between four of 8 U-V gene pairs studied is only around 1-3%, but two genes have divergence of almost 7%

for (McDaniel *et al.*, 2013); one of them is long enough to reliably suggest high divergence, either indicating a longer time since recombination stopped between the U and V regions, or a higher mutation rate and/or lesser selective constraint (divergence from the related species is also high for the high U-V divergence gene, consistent with the two latter possibilities). Even the highest silent site divergence currently found suggests, however, that this is not an ancient system (although genes with higher divergence may be discovered when more genes are analysed).

Plants with very small, or no, non-recombining regions.

Genetic mapping (or related methods that can detect such regions even if they are small, such as bulk segregant analysis) has yet to be applied in many plants with genetic sex determination, and they could reveal non-recombining regions in many plants not currently classified as having sex chromosomes. Indeed, a major currently unanswered question is whether the number of plants with sex-linked regions is currently under-estimated. Such studies are, however, limited by marker density, and very small non-recombining regions may be missed due to insufficient marker density. Indeed, in several plants, genetic sex determination has been established, and a SEX locus controlling gender has been mapped, but no fully sex-linked marker has been found. In kiwifruit (*Actinidia chinensis*), for example, mapping with 644 microsatellite markers still failed to detect any fully sex-linked markers. Other species where the recombination status of the SEX locus is currently uncertain include spinach (Khattak *et al.*, 2006), asparagus (Telgmann-Rauber *et al.*, 2007), and *Populus* species (Yin *et al.*, 2008; Pakull *et al.*, 2011). Such species may, of course, truly lack non-recombining SEX regions. They may either not yet have evolved fully sex-linked regions, or may be single gene systems, which can evolve when a new gene takes over control of flower sex determination after dioecy has become established, replacing an existing sex-determining gene (Bull, 1983; van Doorn & Kirkpatrick, 2007; Vuilleumier *et al.*, 2007; Blaser *et al.*, 2013); Figure 1C illustrates such an event. Takeovers are known in several animal taxa, including insects (Wilkins, 1995; Beye *et al.*, 2003) and fish (Ross *et al.*, 2009; Myosho *et al.*, 2012).

To map SEX loci, high throughput methods including RAD-Seq (Baird *et al.*, 2008) or RNA-Seq transcriptome sequencing (Bergero & Charlesworth, 2011; Chibalina & Filatov, 2011; Muyle *et al.*, 2012; Hough *et al.*, 2014) can now generate

large numbers of markers, overcoming the problem of low marker density. Alternatively, given the large resources sometimes available in crop plants, high density linkage maps can be obtained by “target-sequence capture” (Mamanova *et al.*, 2010). In *Fragaria vesca*, for example, a map was made by first obtaining a low coverage genome sequence, then identifying polymorphisms, and then using an enrichment approach to obtain short sequences (200bp) surrounding each polymorphism. This allowed genotyping of 5417 genes in a mapping family (Tennessen *et al.*, 2013). Another recently developed approach that may be helpful in plants ascertains fully sex-linked sequences by searching short read genome sequence data from multiple individuals for k -mers (short sequences of length k) that appear only in one sex (Carvalho & Clark, 2013). In the section on sex-determining genes below, I outline how this approach has ascertained Y-linked sequences in persimmon (*Diospyros lotus*), in the Ebenaceae (Akagi *et al.*, 2014).

A major difference between ancient animal sex chromosomes and the Y-linked regions of the plants just discussed (with the possible exception of *M. polymorpha*) is the minor extent of gene loss in plants (see section on genetic degeneration below). In plants, sex-linked genes therefore cannot be ascertained by genome sequencing, using their lower coverage in the heterogametic sex. Moreover, assembly of short-read sequences will be difficult, due to sequence divergence and accumulation of repetitive sequences, which occur in non-recombining genome regions (Charlesworth *et al.*, 1994), including plant sex chromosomes (Kubat *et al.*, 2014). Assemblies of the human and papaya Y chromosomes involved deep sequencing of single males, avoiding variants that might confuse assembly, and are restricted to non-heterochromatic regions (Hughes *et al.*, 2010; Wang *et al.*, 2012).

Finally, multiple individuals of each sex are needed to determine sex-specific sequences and distinguish fully and partially sex-linked regions. Even with large family sizes, lack of recombinants does not exclude rare recombination. Population genetic approaches can, however, detect recombination in past generations, even if it occurs very rarely. In papaya, for example, the SEX region adjoins a “collinear region” where the sequenced X and Y chromosomes appear to have the same genes in the same order, unlike the older Y-linked strata, whose assembly includes many rearrangements (Wang *et al.*, 2012). Divergence between the single X and Y sequences so far available is low in the collinear region, indicating that recombination must have continued after it had stopped in the two strata defined by the Y region

inversions described above. Sequence differences between the single X- and Y-linked region so far sequenced may merely be variants that happens to be carried on those particular chromosomes, and not sex-linked in the species as a whole, so this region may prove to be partially sex-linked.

An approach that does include multiple individuals is bulk segregant analysis. This has recently been successful in the grape vine (*V. vinifera*), an XY system. The fully sex-linked genome region is only about 150 kb, and includes only a few fully sex-linked genes, so recombination suppression has clearly not yet extended across any substantial genome region (Picq *et al.*, 2014), consistent with the lack of chromosomal heteromorphism.

The observation that much of the chromosome carrying the SEX locus recombines does not necessarily mean that it is a young system. Instead, it may be an ancient system whose non-recombining region has remained limited to just a part of the chromosome. This is the case in the Ratite birds (Pigozzi, 2011; Vicoso *et al.*, 2013; Zhou *et al.*, 2014). In *Vitis*, most wild species are dioecious, suggesting that dioecy is ancestral, so this could be a plant example of an old-established system that has not evolved recombination suppression. However, divergence between *Vitis* sex-linked genes has not yet been estimated, so the time when the sex-determining region evolved is not yet known.

Comparative genetic mapping

Genetic mapping is also important for detecting differences in the chromosome carrying the SEX region. This can occur when genetic sex-determination evolves independently (or re-evolves after loss of dioecy). For example, genes that are sex-linked in *S. latifolia* do not show sex-linkage in *Silene otites* and *S. colpothpylla* (Mrackova *et al.*, 2008). Phylogenetic analyses of sex systems in *Silene* (Mrackova *et al.*, 2008; Marais *et al.*, 2011) suggest that these species do not have a dioecious common ancestor, so dioecy probably evolved *de novo* in the two lineages, and involved different genes.

Events in which a new gene takes over control of gender, can also cause the SEX loci of related species to be on non-homologous chromosomes (Figure 1C), or to a new location on the same chromosome, as in some animal cases of takeovers (Uno *et al.*, 2008). In *Populus*, as in papaya, the SEX loci appear to be within

pericentromeric regions (Geraldes *et al.*, 2011), but, although these regions map to the same linkage group, their locations differ greatly in the physical maps of different *Populus* species (Yin *et al.*, 2008; Pakull *et al.*, 2011). If confirmed, this suggests a takeover event by a new sex-determining gene on the same chromosome. Independent evolution of separate sexes in different *Populus* lineages is not yet excluded, however, even though almost all Salicaceae are dioecious. Data on the ages of the systems, phylogenetic analysis, and genome sequencing, should help to distinguish between the possibilities. Takeovers or independent evolution both predict that different species should have different sets of genes at their SEX loci, unlike a chromosome rearrangement. Independent evolution predicts that the times since recombination stopped should differ (though this might not be detectable if all species have young systems), while takeover events generating single-locus systems may not have been followed by recombination suppression in the surrounding genome region.

Why does suppressed recombination evolve?

The repeated evolution of regions without crossing over between sex chromosomes strongly suggests a causal connection with the evolution of sex-determining regions (only the centromeric and pericentromeric regions of autosomes generally have suppressed crossing over). The evolutionary strata of sex chromosomes discussed above prove that suppressed recombination often evolves after a sex-determining system is established. Some disadvantage to recombinant genotypes must clearly be involved. Such situations probably occur both during the initial evolution of dioecy, and also later, as males and females evolve in the absence of constraints imposed by the other sex functions. Briefly, as illustrated in Figure 2, separate sexes in plants probably often evolved from hermaphroditic or monoecious ancestors, often called “cosexual” species (Lloyd, 1982).

The change from cosexuality to dioecy probably involves a mutation creating females (a male-sterility mutation in an initially hermaphroditic species, or a mutation suppressing some or all female flowers in an initially monoecious species, or replacing them with male flowers), and then one or more female-suppressing mutations, creating males or male-biased plants (Westergaard, 1958; Charlesworth & Charlesworth, 1978). (I can find no cases where dioecy in plants evolved from environmental sex determination, though this seems possible in principle).

[Figure 2 about here]

In this scenario, male-promoting mutations (suppressing femaleness) clearly reduce females' fitness, and are therefore most likely to spread if linked to the gene causing femaleness, which minimises the conflict. If a two-gene polymorphism results, selection against recombinants will generate linkage disequilibrium (with the X associated with male-sterility and the Y with female-suppressor alleles, see Figure 2). Suppressed recombination is therefore favoured, and may evolve (Charlesworth & Charlesworth, 1980; Bull, 1983), creating a male-determining Y chromosome.

If however, a male-specific female-suppressing mutation occurs, no harm is caused to females; if sufficiently advantageous in males, such a mutation can spread, even if unlinked to the femaleness gene, yielding a single gene sex-determining system (Figure 2A), and no selection for closer linkage with the gene causing femaleness (Muller, 1932; Charlesworth & Charlesworth, 1978).

Much genetic evidence supports the two-gene model for plant sex-determination, rather than one with sex-specific gene actions. First, in several species (or intercrosses of dioecious plants with close non-dioecious relatives) three allelic types at the sex-determining locus control whether individuals are (i) females, (ii) males, or (iii) hermaphrodites or monoecious functional hermaphrodites (Westergaard, 1958). Second, in papaya and grape vine, humans have selected individuals that have Y-linked regions that do not suppress female functions (Wang *et al.*, 2012; Picq *et al.*, 2014). Similarly, in *Silene latifolia*, deletions detectable through loss of Y-linked markers can create hermaphrodites and neuter plants (Fujita *et al.*, 2011 and references therein). These plants' Y-linked regions must therefore carry suppressors of female functions whose loss does not affect male functions, and distinct maleness factor(s) elsewhere on the chromosome. Thirdly, in the strawberry species *Fragaria virginiana*, two closely, but not completely, linked genes with the expected phenotypes have been found (Spigler *et al.*, 2008), while a related *Fragaria* species has suppressed recombination (Goldberg *et al.*, 2010). These species' sex-determining regions probably evolved independently (Goldberg *et al.*, 2010), but the results nevertheless suggest recombination suppression evolving in response to a two-locus polymorphism.

376 The evolution of dioecy probably often involves further sexually antagonistic
 377 mutations, leading to further selection to suppress recombination, and potentially
 378 generating younger strata. For example, dioecy has often evolved from monoecy
 379 (Renner, 2014), and full maleness may involve successive increases in the proportion
 380 of investment in male flowers (Figure 2B), each involving sexually antagonistic
 381 “trade-offs”, because each must decrease the proportion invested in female flowers.
 382 Variable degrees of maleness are indeed seen in the monocotyledon *Sagittaria*
 383 *latifolia*, in the Alismataceae (Dorken & Barrett, 2004), *Spinacia oleracea* in the
 384 Chenopodiaceae (Onodera *et al.*, 2011) and *Urtica dioica* (Glawe & Jong, 2009).
 385 Similarly, when the ancestral state is hermaphroditism, evolution of dioecy often
 386 involves “inconstant males” with partial female function (for example, producing
 387 some fruits in favourable conditions). Species where genetic variation in male
 388 functions seems likely include *Antennaria dioica* in the Asteraceae (Ubisch, 1936)
 389 and *Euonymus europaeus* in the Celastraceae (Webb, 1979), but these have not been
 390 investigated using genetic markers to map the factors. Even after complete
 391 unisexuality has evolved, male and female functioning may be sub-optimal, and
 392 improvements to each sex may often reduce functions of the other. In *S. latifolia*, for
 393 example, female fecundity is enhanced by making large flowers, but fertility is
 394 highest for males with many small flowers (Delph & Herlihy, 2012). Just as outlined
 395 above for sterility mutations, a mutation benefitting one sex at the expense of the
 396 other is most likely to invade but not spread throughout the population; if such a
 397 polymorphism is established, it creates selection for reduced recombination with the
 398 sex-determining locus, if it arises at a locus closely linked to the SEX locus (Rice,
 399 1987; Jordan & Charlesworth, 2012).

400 Testing for the trade-offs and conflicts assumed in these scenarios, and for
 401 involvement of sexually antagonistic polymorphisms in the PAR regions of sex
 402 chromosomes is clearly a major task for future work. An approach that can potentially
 403 detect sexually antagonistic variation is QTL analysis within the two sexes separately,
 404 as proposed and implemented in *Silene latifolia* (Scotti & Delph, 2006; Delph *et al.*,
 405 2010). This detected several autosomal and PAR QTLs, and, interestingly, the latter
 406 appeared only in the analysis of males, implying that their phenotypic effects are not
 407 expressed in females. Such male-specific expression is consistent with a past conflict
 408 between the sexes that has been resolved in later evolution, as seems to have occurred
 409 for some sexually selected male coloration genes in the PAR of a fish, the guppy,

Poecilia reticulata (Lindholm & Breden, 2002). Male benefit alleles with male-specific expression no longer harm females, and will spread throughout the population; some other selection is therefore required to maintain the QTL variation, perhaps environmental differences (Scotti & Delph, 2006). In *S. latifolia*, for example, thin leaves appear to be disadvantageous to males only in dry years (Delph *et al.*, 2011). The *S. latifolia* QTL analysis used dominant AFLP markers, but codominant markers now available in this plant's PAR, and obtainable in other plants, will permit future analyses of variation in natural populations. This may detect factors whose conflict has not been resolved, corresponding to the situation that creates selection for reduced recombination in the theoretical models of sexually antagonistic PAR genes.

Recombination suppression: mechanisms

Non-recombining regions may eventually evolve to encompass a large region of the chromosome carrying the sex-determining loci or locus. Studies of young plant sex chromosomes may be valuable for studying the mechanistic basis of recombination suppression, and whether it generally involves infrequent, large-scale events like inversions, or smaller shifts in the position of the PAR boundary.

If chromosome inversions cause recombination suppression in SEX regions (Lahn & Page, 1999), the region will often include many non-sex-determining genes. In papaya, two Y chromosome inversions indeed seem to be involved (including 10 genes with both X and Y copies in the older stratum includes, and 16 in the newer one, Wang *et al.*, 2012). In closely related dioecious *Vasconcellea* species, alleles of several papaya fully sex-linked genes are not associated with gender (Gschwend *et al.*, 2011). Unlike the *Silene* situation described above, this probably does not reflect independent evolution of dioecy in *Carica* and *Vasconcellea*, as BAC-FISH experiments found sex-linked regions including several homologous sequences in similar locations on the largest chromosome of both species (Iovene *et al.*, 2015). Recombination suppression has therefore probably remained restricted in *V. parviflora* to a genome region near the SEX locus, whereas it has spread across a wider region in *C. papaya*. This is testable by sequencing to ask whether *V. parviflora* genes homologous to *C. papaya* genes in the older sex-linked stratum have distinct X and Y haplotypes like those of papaya. The alternative that the *V. parviflora* long arm has become a new recombination-suppressed stratum seems unlikely, because the

443 chromosomal positions of all nine relevant BACs in an outgroup, *Jacaratia spinosa*,
444 were found to be similar to those in *V. parviflora*, so the inversions probably occurred
445 in the *C. papaya* lineage.

446 Recombination suppression mechanisms other than inversions may, however,
447 exist, including modifiers controlling the number of crossover events, restricting them
448 to certain genome regions, or restricting crossing over to only one sex. In one of the
449 two human PARs, for example, crossovers are localised very differently in male and
450 female meiosis (Hinch *et al.*, 2014). Some young sex chromosome systems may still
451 be in the process of undergoing recombination suppression. If recombination varies
452 between individuals, or between closely related species that can be interbred, genetic
453 studies can potentially identify the factors involved. In some populations of frog
454 species, male-specificity of microsatellite alleles differs between populations,
455 implying that the XY pair shows suppressed recombination only in some populations
456 (Dufresnes *et al.*, 2014). Apparently similar variation was inferred for an anonymous
457 sequence marker within the plant species *Bryonia dioica* (Oyama *et al.*, 2009), which
458 should be studied further. In *Silene latifolia*, recombination suppression appears to
459 vary between families for several genes (Bergero *et al.*, 2013).

460

461 **Old-established sex chromosome systems**

462 Old sex chromosomes also exist among plants, for example among liverworts
463 (Okada *et al.*, 2001), but have been less studied than young plant systems. As
464 molecular approaches and phylogenetic analyses are extended to studying more plant
465 sex chromosome systems, it will be interesting to include taxa with high proportions
466 of dioecious species, such as the palm, Vitaceae and Ebenaceae (including *Diospyros*
467 *lotus*, see below) families, to test whether dioecy is ancestral and estimate the time
468 when recombination stopped, or has evolved several times (as may be the case in the
469 Salicaceae discussed above).

470 Old systems are particularly interesting for investigating genetic degeneration
471 and repetitive sequence accumulation, which occur over large evolutionary
472 timescales. The potentially large range of ages of dioecious plant sex chromosome
473 systems will allow the time-course of sex chromosome evolution to be studied. Old
474 plant systems may also help us understand why recombination suppression sometimes
475 fails to evolve.

The evidence for old-established systems is currently incomplete, and age estimates based on sequence divergence are lacking. There are currently no dense genetic maps for plants that seem likely to have old XY systems, and, so far, genetic mapping in these systems has largely used non-genic markers such as AFLPs and microsatellites. These are excellent for testing for a non-recombining (sex chromosome-like) region, determining which is the heterozygous sex, and estimating the proportion of the chromosome that is fully sex-linked. However, as explained above, estimating the age of a sex chromosome system, and the time when recombination stopped, requires X-Y sequence divergence estimates, based on ascertaining sex-linked genes and sequencing them.

In the absence of divergence data, the observation that a sex chromosome system is heterochromatic and heteromorphic might be thought to suggest that it is old-established, especially in plant families that include distantly related dioecious species, such as date palms (Al-Mahmoud *et al.*, 2012). For example, *Rumex acetosa* belongs to a clade that may have been dioecious for 15–16 MYA (Navajas-Pérez *et al.*, 2005), but X-Y divergence has not been estimated. Its Y chromosomes are heterochromatic (Shibata *et al.*, 2000; Mariotti *et al.*, 2008), unlike those of other cytologically well-studied plants such as *Silene latifolia* and *S. dioica* (Grabowska-Joachimciak & Joachimciak, 2002; Kubat *et al.*, 2014), which are estimated to be younger (see above). However, heterochromatin can evolve rapidly, as in papaya. Another example is *Coccinia grandis*, within a wholly dioecious genus of 27 species (in the Cucurbitaceae, another family with many dioecious species, often with XY heteromorphism). Its male genome C-value is 10% larger than that of females, indicating that the Y chromosome is much larger than the X, and the entire Y is heterochromatic (Sousa *et al.*, 2012), yet phylogenetic analysis suggests that these characteristics evolved recently (Holstein & Renner, 2011).

Sex chromosome heteromorphism can also arise in young systems, for example through fusions with autosomes, as in *Rumex hastatulus* (Smith, 1964) and possibly spinach (Araratjan, 1939). The systems in *Cannabis sativa* (Peil *et al.*, 2003; Sakamoto *et al.*, 2005) and *Humulus lupulus* (hops) in the Cannabaceae, whose Y chromosome is heterochromatic (Westergaard, 1958) are probably much older.

Studies of old systems are also needed to test the prediction that other sex-determining systems are derived from XY systems (Charlesworth & Charlesworth, 1978). Again plants may be very helpful, as systems with male-determining Y

chromosomes probably evolve first, as outlined above, but ZW systems also exist, and it can be tested whether the frequencies of such systems increases over time. X-autosome balance systems are also probably derived from XY systems (and potentially allow loss of the Y chromosome, and evolution of an X0 male genotype). However, it has again not yet been demonstrated that such species tend to be older than other plant sex-determining systems. Absence of carpel development in males or stamen development in females, as in hops, may also indicate an ancient system (but might simply be due to a long history of unisexual flowers, for example because dioecy has evolved from monoecy); so far, only one fully sex-linked genetic marker locus has been found in hops (Jakse *et al.*, 2008).

Genetic degeneration: the need for empirical data in a phylogenetic setting

Ancient systems are also of great interest for studying genetic degeneration (gene loss or loss of function). In diploid organisms, only the Y chromosomes are predicted to degenerate, because X chromosomes recombine in the XX females, whereas Y-linked regions do not, and are subject to several processes that allow detrimental mutations to increase in frequency in the population of Y-linked alleles, or even to become fixed in this population, as recently reviewed (Bachtrog, 2008). In haploid plants, however, the complete lack of recombination across the entire sex-linked region predicts similar degeneration of both U and V chromosomes (Bull, 1983). Genes affecting non-sex functions should not degenerate or become lost, so the female-determining U region should lose only male function genes, and the male-determining V region only female function genes (Figure 3C).

Figure 3 about here

Haploid plants with separate sexes of gametophytes are ideal for studying this prediction. In *Marchantia polymorpha*, a species whose sex chromosomes carry highly diverged sequences, the V has been studied in detail, but analysis of the U chromosome is currently incomplete (Okada *et al.*, 2001). In the brown alga, *Ectocarpus siliculosus*, however, about 24 genes were found in the fully sex-linked regions (either the U or V regions, or both), of which 7 were not detected in the V and 9 in the U (Ahmed *et al.*, 2014). This is in apparent agreement with Bull's prediction;

however, without an outgroup, gene movements onto one sex chromosome, but not the other, cannot be excluded. To determine whether suitably close non-dioecious relatives exist (and avoid species that might have reverted from dioecy to a non-dioecious state), phylogenetic relationships of the species must be known. This is often difficult for closely related species, a frequent situation relevant to the evolution of sex chromosomes. Nevertheless, among plants, sets of species should exist with good phylogenies well suited for future work estimating ancestral character states, and changes in states, (Maddison & Leduc-Robert, 2013).

Diploid dioecious plants also have extended haploid life cycle stages, which may also cause genetic degeneration of non-recombining sex chromosomal regions to be minor (Figure 3B). Around 2/3 of plant genes are expressed in male gametophytes of angiosperms (Tanksley *et al.*, 1981; Gorla *et al.*, 1986; Honys & Twell, 2003). Therefore, only genes with no important pollen functions should be lost from plant SEX regions, or lose their functions; the limited evidence so far about loss of genes from the *S. latifolia* Y chromosome is consistent with this expectation (Guttman & Charlesworth, 1998; Chibalina & Filatov, 2011). Degeneration might be thus restricted to around 1/3 of genes (or possibly somewhat higher, if expression of some pollen-expressed genes is not important, and purifying selection maintaining their functions is consequently weak). The few current estimates, from the unrelated plants *S. latifolia* and *Rumex hastatulus*, suggest that fewer than 30% of Y-linked genes have lost expression (Bergero & Charlesworth, 2011; Chibalina & Filatov, 2011; Hough *et al.*, 2014). In contrast, such regions are almost completely degenerated in the best studied animals, such as species of *Drosophila* (Muller, 1950), mammals (Skaletsky *et al.*, 2003) and those birds that have extensive fully W-linked regions (Zhou *et al.*, 2014), and possibly in part of the much younger Y chromosome of the threespine stickleback (Ross & Peichel, 2008; Yoshida *et al.*, 2014). Large genome regions that stopped recombining recently and carry many genes driving the degeneration processes, such as the neo-Y chromosome of *Drosophila miranda*, have quickly lost functions of large fractions of genes (Bachtrog *et al.*, 2008). However, the regions of the two plants so far studied that recently became fully sex-linked probably include many fewer genes than the *D. miranda* region, so that the small extent of gene losses in these young systems is not surprising. It will be interesting to study older plant systems.

Genetic degeneration in young plant sex chromosomes, and in young evolutionary strata in older systems, is also of interest. The first step after recombination is suppressed between Y- and X-linked regions may be accumulation of repetitive sequences, including transposable elements. Such insertions may decrease expression of Y-linked alleles, even before mutations in the coding regions, or in non-coding regions that control the gene's expression. This appears to be the case in *Drosophila albomicans* (Zhou & Bachtrog, 2012).

However, plants with sex-determining loci within rarely recombining pericentromeric regions, such as papaya and *Populus* species, are not well suited for studying genetic degeneration, because accumulation of maladaptive sequence changes and of repetitive sequences are also expected in pericentromeric genome regions (Charlesworth *et al.*, 1986). It will therefore be difficult to detect extra effects of the evolution of sex-determining genes in the region. For example, in papaya, gene density is low in the sex-linked region, but this is not wholly due to loss of genes; accumulation of repeated sequences has also reduced gene density (Wang *et al.*, 2012).

Dosage compensation

In sex chromosome systems where Y-linked gene expression is reduced, or Y-linked genes have been lost, dosage compensation has sometimes evolved, and it is therefore interesting to test whether X-linked alleles of plant genes whose Y-linked copies have lost function are expressed at higher levels in males than females. There is currently no clear evidence that this occurs in *Silene latifolia* or *Rumex hastatulus*, but partial compensation cannot yet be excluded (Chibalina & Filatov, 2011; Muyle *et al.*, 2012; Hough *et al.*, 2014; Bergero *et al.*, 2015).

Plant sex-determining loci

To identify sex-linked regions and determine whether males or females are the heterozygous sex, it is not necessary to find the gene(s) controlling male or female development. As explained above, it suffices to find genetic markers, even anonymous ones, such as AFLPs, that co-segregate with sex. However, plant sex-determining loci are interesting in several ways, including for identifying the hypothesised two or more genes causing male- and female-sterility during the evolution of dioecy. If sex-determining genes can be discovered, sequence divergence

between their sex-linked alleles may also help estimate the time when recombination first stopped. With the possibility of dense marker development and genome sequences, renewed efforts are being made to identify plant sex-determining genes, and progress can be expected in the next few years.

The approach of testing known flower development genes has been largely superseded by high-throughput sequencing methods. Searches have found MADs-box and ABC(DE) genes involved in flower whorl development on the sex chromosomes of *Silene latifolia* and *Asparagus officinalis* (Matsunaga *et al.*, 2003; Park *et al.*, 2003; Cegan *et al.*, 2010; Nishiyama *et al.*, 2010; Penny *et al.*, 2011). However, genes that control floral organ identity are not generally promising candidates. They might be involved in species with complete absence of one sex organs (“Type I” of Mitchell & Diggle, 2005; Ramos *et al.*, 2014). In many dioecious plants, however, both male and female floral organs are initiated in flowers of both sexes, and the development of opposite sex organs is later interrupted.

Alternative approaches also encounter difficulties due to the numerous candidates whose loss of function can produce male or female sterility. For example, as mentioned above, deletion mapping of the *S. latifolia* Y chromosome has established that separate loci exist whose deletion causes abortion or incomplete development of stamens, or removes the suppression of pistils that occurs in wild-type males, creating hermaphrodite flowers (Farbos *et al.*, 1999; Lardon *et al.*, 1999; Zluvova *et al.*, 2005; Bergero *et al.*, 2008; Fujita *et al.*, 2011). However, these deletions probably involve loss of many fully sex-linked genes other than the ones causing these phenotypes, and this is supported by the observation that pollen carrying deleted Y chromosome regions often has low ability to fertilise ovules (Lardon *et al.*, 1999). When the sex-linked region is large, it will be difficult to identify the genes responsible for the evolution of dioecy unless small deletions can be generated and identified using dense mapping of sequences lost from deleted genotypes.

This problem also hinders attempts to identify genes involved in gender determination using mutations, including mutations induced by EMS or irradiation (Ohnishi, 1985; Christensen *et al.*, 1998; Honys & Twell, 2004; Wellmer *et al.*, 2006; Chang *et al.*, 2011), or by studying genes with different expression in flower buds of the two sexes. Moreover, many genes have stamen- or pistil-specific expression, and will be non-expressed in buds of one sex purely because the relevant structures are

absent. Distinguishing such downstream acting genes from the sex-determiners themselves requires establishing sex linkage. If, however, the fully sex-linked region includes many genes, the problem of having too many candidates with suitable function is not eliminated. In addition, expression differences may not be involved (for example, male sterility can involve mutations in coding sequences, and the mutant alleles may be present in mRNA).

Small sex-linked regions may offer the best prospects for identifying the sex-determining genes, because fewer candidates need to be considered. A candidate Y-linked gene has been proposed in persimmon (Akagi *et al.*, 2014). This study started by identifying sex-linked genes, using pools of males and females from a full-sib family, and their sex-linkage was confirmed in samples of unrelated males and females. Efforts were made to ensure that most Y-linked genes present in transcripts were detected, by employing RNA-Seq, and 22 expressed sequences were identified. The total length of sex-specific sequences was only 1Mb, suggesting a small fully sex-linked region. One candidate for involvement in sex determination was found. This gene (named *OGI*) is expressed only in male flower buds. *OGI* is a duplication onto the Y-linked region of an autosomal gene called *MeGI* that expresses a male-suppressing regulatory RNA in females. Because no X copy exists, X-Y divergence cannot be estimated, but divergence from the presumed autosomal progenitor is high, and Y-linked *OGI* sequences were detected in other species of Ebenaceae, suggesting an old-established Y-linked duplication. Low divergence was found between the X- and Y-linked alleles of other sex-linked genes (silent site divergence of 12 XY allele pairs was below 2%), suggesting that a younger stratum evolved recently.

The proposed scenario for sex-determination in persimmon is that the Y-linked *OGI* gene opposes *MeGI*'s male-suppressing action. This form of gene action that could act in the heterozygous state, and should increase male functions, and the processes in the two sexes may indeed conflict, as proposed for the female suppressor in the two-gene model outlined in Figure 2A above. It is currently unclear how females evolved. The *MeGI* male suppressing factor is autosomal, and is therefore unlikely to represent the male-sterility gene in the two-gene model. *OGI* could therefore be an example of a single-locus sex-determining gene that evolved by a take-over event, if searches fail to find a femaleness factor.

Although, as mentioned already, reversals and re-evolution of dioecy can complicate comparative studies and hinder inferences of the ages of the origins of

677 dioecy, they may also be very helpful in revealing the genetic basis of dioecy
678 (Westergaard, 1958), and molecular studies of such hermaphrodites could help
679 identify plant sex-determining genes. In papaya and *Vitis*, hermaphrodites are
680 commercially successful crop plants. The Y chromosomes in these hermaphrodites do
681 not suppress female functions, but their sequences are very similar to those of males
682 (Picq *et al.*, 2014; Van Buren *et al.*, 2015), and they probably have no large deletions,
683 making them ideal for identifying the gene whose loss causes reversion to
684 hermaphroditism, a good candidate for the female suppressor involved in the
685 evolution of dioecy.

686 The hypothesised X-linked genes responsible for the male sterility of females in
687 dioecious plant species are likely to be even harder to identify, but this may be
688 possible in systems where suppressed recombination has not yet evolved. If two
689 incompletely linked sex determining genes exist, hermaphrodite recombinants, and
690 recombinants with the male sterility allele of females and the female suppressor of
691 males, should arise. With the modern ability to identify the region, and genotype
692 closely linked markers, as in *Fragaria* species (Tennessen *et al.*, 2013), it should be
693 possible to check that these phenotypes are indeed associated with recombination, and
694 to pinpoint both genes.

695 Once the genes are identified in some plant species, this will open the way for
696 testing whether the same genes are sex-linked in other dioecious plants. Given that
697 large numbers of genes affect flower and inflorescence development, different genes
698 may be involved in different angiosperm lineages, rather than the same genes being
699 repeatedly involved. If so, plants will differ from major animal groups such as insects,
700 which share sex-determination pathways across major taxa (Saccone *et al.*, 2002;
701 Beye *et al.*, 2003; Pomiankowski *et al.*, 2004; Pane *et al.*, 2005). In plants, sterility
702 factors may have to be identified, and their actions investigated, in individual genera
703 and species. Moreover, it should not be assumed that the sex-determining genes
704 necessarily function during flower development, or cause sterility. In monoecious
705 plants, a state that is ancestral to many dioecious species (Renner & Ricklefs, 1995;
706 Renner & Won, 2001), they might instead control the proportions of male and female
707 flowers, perhaps at developmental stages before flower parts are initiated (Figure 2B).
708 Unisexuality may be much more ancient than dioecy, and early, complete abortion of
709 male or female parts may be ancestral.

710

Conclusions

It is now technically feasible to use young sex chromosomes in non-model plants to test hypotheses about the initial evolution of suppressed recombination, and to study the time course of later evolution of sex chromosomes in older systems, as has been initiated in some animal systems (Bachtrog *et al.*, 2009). Young plant sex-linked systems should also be suitable for studying the earliest adaptations to dioecy, which have so far been little studied. The change from cosexuality to unisexuality may be accompanied by considerable expression changes, if unisexuals are released from conflicts between the two sex functions, so that changes can occur to optimise expression in each sex. For example, the non-dioecious *S. vulgaris* appears suitable as an outgroup for studying the evolution of changes in expression in the dioecious species *S. latifolia* (Marais *et al.*, 2011). Because hermaphrodite *S. vulgaris* individuals have both stamens and pistils, differences in unisexual individuals of the dioecious species that are caused directly by loss of these structures should be distinguishable from changes in expression of genes that are expressed in non-sex-specific structures. Such studies can potentially discover genes that can be expressed in both sexes, but that change when dioecy evolves, and evaluate whether, as has been predicted, the sex chromosomes, including the PAR, carry unexpectedly large numbers of such genes (Vicoso & Charlesworth, 2006; Vicoso *et al.*, 2013).

Acknowledgements: I thank S.C.H. Barrett, S.I Wright and J. Leebens-Mack for comments on an earlier version of this manuscript.

References

- Ahmed S, Cock JM, Pessia E, Luthringer R, Cormier A, Robuchon M, Sterck L, Peters AF, Dittami SM, Corre E, Valero M, Aury J, Roze D, Peer YVd, Bothwell J, Marais GAB, Coelho SM. 2014. A haploid system of sex determination in the brown alga *Ectocarpus* sp. *Current Biology* **24**(17): 1945–1957.
- Akagi T, Henry IM, Tao R, Comai L. 2014. A Y-chromosome–encoded small RNA acts as a sex determinant in persimmons. *Science* **346**: 646–650.
- Al-Mahmoud M, Al-Dous E, Al-Azwani E, Malek J. 2012. DNA-based assays to distinguish date palm (Arecaceae) gender. *American Journal of Botany* **99**(1): E7–E10.
- Alstrom-Rapaport C, Lascoux M, Wang YC, Roberts G, G.A T. 1998. Identification of a RAPD marker linked to sex determination in the basket willow (*Salix viminalis* L.). *Journal of Heredity* **89**: 44–49.

- Araratjan A. 1939.** Heterochromosomes in the wild spinach. *Comptes Rendus (Doklady) Acad. Sci. URSS* **24**: 56-57.
- Bachtrog D. 2008.** The temporal dynamics of processes underlying Y chromosome degeneration. *Genetics* **179**(3): 1513-1525.
- Bachtrog D. 2012.** Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nature Reviews Genetics* **14**(2): 113 -124.
- Bachtrog D, Hom E, Wong K, Maside X, Jong Pd. 2008.** Genomic degradation of a young Y chromosome in *Drosophila miranda*. *Genome Biology* **9**(2): R30.
- Bachtrog D, Jensen J, Zhang Z. 2009.** Accelerated adaptive evolution on a newly formed X chromosome. *PLoS Biology* **7**(4): e1000082.
- Bachtrog D, Kirkpatrick M, Mank J, McDaniel SF, Pires JC, Rice WR, Valenzuela N. 2011.** Are all sex chromosomes created equal? *Trends in Genetics* **27**(9): 350-357.
- Baird N, Etter P, Atwood T, Currey M, Shiver A, Lewis Z, Selker E, Cresko W, Johnson E. 2008.** Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLOS ONE* **3**(10): e3376.
- Baker K, Bayer M, Cook N, Dressig S, Dhillon T, J Russell, Hedley P, Morris J, Ramsay L, Colas I, Waugh R, Steffenson B, Milne I, Stephen G, D Marshall, Flavell A. 2014.** The low-recombining pericentromeric region of barley restricts gene diversity and evolution but not gene expression. *Plant Journal* **79**(6): 981-992.
- Bellott D, Hughes JF, H Skaletsky, Brown LG, Pyntikova T, Cho T, Koutseva N, S Zaghlul, Graves T, Rock S, Kremitzki C, Fulton RS, Dugan S, Ding Y, Morton D, Khan Z, Lewis L, Buhay C, Wang Q, Watt J, Holder M, Lee S, Nazareth L, Rozen S, Muzny DM, Warren WC, Gibbs RA, R.K. Wilson, Page DC. 2014.** Mammalian Y chromosomes retain widely expressed dosage-sensitive regulators *Nature* **508**: 494-499.
- Bergero R, Charlesworth D. 2011.** Preservation of the Y transcriptome in a 10MY old plant sex chromosome system. *Current Biology* **21**(12): 1470-1474.
- Bergero R, Charlesworth D, Filatov D, Moore R. 2008.** Defining regions and rearrangements of the *Silene latifolia* Y chromosome. *Genetics* **178**: 2045-2053.
- Bergero R, Forrest A, Kamau E, Charlesworth D. 2007.** Evolutionary strata on the X chromosomes of the dioecious plant *Silene latifolia*: evidence from new sex-linked genes. *Genetics* **175**: 1945-1954.
- Bergero R, Qiu S, Charlesworth D. 2015.** Gene loss from a plant sex chromosome system *Current Biology* **25**: in press (doi.org/10.1016/j.cub.2015.03.015).
- Bergero R, Qiu S, Forrest A, Borthwick H, Charlesworth D. 2013.** Expansion of the pseudoautosomal region and ongoing recombination suppression in the *Silene latifolia* sex chromosomes. *Genetics* **194**(3): 673-686.
- Beye M, Hasselmann M, Fondrk MK, Page RE, Omholt SW. 2003.** The gene *csd* is the primary signal for sexual development in the honeybee and encodes an SR-type protein. *Cell* **114**: 419-429.
- Blaser O, Grossen C, Neuenschwander S, Perrin N. 2013.** Sex-chromosome turnovers induced by deleterious mutation load. *Evolution* **67**(3): 635-645.
- Bull JJ. 1983.** *Evolution of Sex Determining Mechanisms*. Menlo Park, CA: Benjamin/Cummings.
- Carvalho A, Clark AG. 2013.** Efficient identification of Y chromosome sequences in the human and *Drosophila* genomes. *Genome Research* **23**: 1894-1907.

797 **Cegan R, Marais G, Kubekova H, Blavet N, Widmer A, Vyskot B, Dolezel J,**
798 **Safar J, Hobza R. 2010.** Structure and evolution of *Apetala3*, a sex-linked
799 gene in *Silene latifolia*. *BMC Plant Biology* **10**: 1.

800 **Chang F, Wang Y, Wang S, Ma H. 2011.** Molecular control of microsporogenesis in
801 *Arabidopsis*. *Current Opinion in Plant Biology* **14**: 66–73.

802 **Charlesworth B, Charlesworth D. 1978.** A model for the evolution of dioecy and
803 gynodioecy. *Amer. Nat.* **112**: 975-997.

804 **Charlesworth B, Sniegowski P, Stephan W. 1994.** The evolutionary dynamics of
805 repetitive DNA in eukaryotes. *Nature* **371**: 215-220.

806 **Charlesworth B, Stephan W, Langley CH. 1986.** The evolution of restricted
807 recombination and the accumulation of repeated DNA sequences. *Genetics*
808 **112**(4): 947-962.

809 **Charlesworth D 1985.** Distribution of dioecy and self-incompatibility in
810 angiosperms. In: Greenwood PJ, Slatkin M eds. *Evolution - Essays in Honour*
811 *of John Maynard Smith*. Cambridge: Cambridge University Press, 237-268.

812 **Charlesworth D, Charlesworth B. 1980.** Sex differences in fitness and selection for
813 centric fusions between sex-chromosomes and autosomes. *Genetical Research*
814 *(Cambridge)* **35**: 205-214.

815 **Cherif E, Zehdi A, Castello K, Chabrilange N, Abdoukader S, Santoni S, Sahli-**
816 **Hannachi A, Glemin S, Aberlenc-Bertossi F. 2012.** Male specific DNA
817 markers provide genetic evidence of an XY chromosome system and
818 recombination arrest and allow to trace paternal lineages in date palm. *New*
819 *Phytologist* **197**(2): 409–415.

820 **Chibalina M, Filatov D. 2011.** Plant Y chromosome degeneration is retarded by
821 haploid purifying selection. *Current Biology* **21**(12): 1475–1479.

822 **Christensen C, Subramanian S, Drews G. 1998.** Identification of gametophytic
823 mutations affecting female gametophyte development in *Arabidopsis*.
824 *Developmental Biology* **202**(1): 136-151.

825 **Cortez D, Marin R, Toledo-Flores D, Froidevaux L, Liechti A, Waters PD,**
826 **Grützner F, Kaessmann H. 2014.** Origins and functional evolution of Y
827 chromosomes across mammals. *Nature* **508**: 488–493.

828 **Delph L, Arntz A, Scotti-Saintagne C, Scotti I. 2010.** The genomic architecture of
829 sexual dimorphism in the dioecious plant *Silene latifolia*. *Evolution* **64**(10):
830 2873-2886.

831 **Delph LF, Andicoechea J, Steven J, Herlihy C, Scarpino S, Bell D. 2011.**
832 Environment-dependent intralocus sexual conflict in a dioecious plant *New*
833 *Phytologist* **192**(2): 542-552.

834 **Delph LF, Herlihy CR. 2012.** Sexual, fecundity, and viability selection on flower
835 size and number in a sexually dimorphic plant. *Evolution* **66**(4): 1154–1166.

836 **Dorken ME, Barrett SCG. 2004.** Sex determination and the evolution of dioecy
837 from monoecy in *Sagittaria latifolia* (Alismataceae). *Proceedings of the Royal*
838 *Society of London B.* **271**: 213-219.

839 **Dufresnes C, Stöck M, Brelsford A, Perrin N. 2014.** Range-wide sex chromosome
840 sequence similarity supports occasional XY recombination in European tree
841 frogs (*Hyla arborea*). *PLOS ONE* **9**(6): e97959.

842 **Farbos I, Veuskens J, Vyskot B, Oliveira M, Hinnisdaels S, Aghmir A, Mouras**
843 **A, Negrutiu I. 1999.** Sexual dimorphism in white campion: complex deletion
844 on the Y chromosome results in a floral asexual type. *Genetics* **151**: 1187-
845 1196.

- Filatov D, Howell EC, Groutides C, Armstrong S. 2008.** Recent spread of a retrotransposon in the *Silene latifolia* genome, apart from the Y chromosome. *Genetics* **181**: 811 - 817.
- Fujita N, Torii C, Ishii K, Aonuma W, Shimizu Y, Kazama Y, Abe T, Kawano S. 2011.** Narrowing down the mapping of plant sex-determination regions using new Y chromosome-specific markers and heavy-ion beam irradiation-induced Y deletion mutants in *Silene latifolia*. *G3* **2**(2): 271-278.
- Geraldes A, Pang J, Thiessen N, Cezard T, Moore R, Zhao Y, Tam A, Wang S, Friedmann M, Birol I, Jones S, Cronk Q, Douglas C. 2011.** SNP discovery in black cottonwood (*Populus trichocarpa*) by population transcriptome resequencing. *Molecular Ecology Resources* **11**: 81-92.
- Glawe GA, Jong TJD. 2009.** Complex sex determination in the stinging nettle *Urtica dioica*. *Evolutionary Ecology* **23**(4): 635-649.
- Goldberg MT, Spigler RB, Ashman TL. 2010.** Comparative genetic mapping points to different sex chromosomes in sibling species of wild strawberry (*Fragaria*). *Genetics* **186**: 1425–1433.
- Gorla MS, Frova G, Binelli G, Ottaviano E. 1986.** The extent of gametophytic-sporophytic gene expression in maize. *Theoret. Appl. Genet.*(42-47).
- Grabowska-Joachimik A, Joachimik A. 2002.** C-banded karyotypes of two *Silene* species with heteromorphic sex chromosomes. *Genome* **45**(2): 243-252.
- Gschwend A, Yu Q, Tong EJ, Zeng F, Han J, Van Buren R, Aryal R, Charlesworth D, Moore PH, Paterson AH, Ming R. 2011.** Rapid divergence and expansion of the X chromosome in papaya. *Proceedings of the National Academy of Sciences of the United States of America*: PNAS MS# 2011-21096.
- Guttman DS, Charlesworth D. 1998.** An X-linked gene has a degenerate Y-linked homologue in the dioecious plant *Silene latifolia*. *Nature* **393**: 263-266.
- Hinch AG, N Altemose, Noor N, Donnelly P, Myers SR. 2014.** Recombination in the human pseudoautosomal region PAR1. *PLoS Genet* **10**: e1004503.
- Holstein N, Renner S. 2011.** A dated phylogeny and collection records reveal repeated biome shifts in the African genus *Coccinia* (Cucurbitaceae). *BMC Evolutionary Biology* **18**(28).
- Honys D, Twell D. 2003.** Comparative analysis of the Arabidopsis pollen transcriptome. *Plant Physiology* **132**(2): 640-652.
- Honys D, Twell D. 2004.** Transcriptome analysis of haploid male gametophyte development in Arabidopsis. *Genome Biology* **5**(11): R85.
- Hough J, Hollister JD, Wang W, Barrett SCH, Otto SP. 2014.** Genetic degeneration of old and young Y chromosomes in the flowering plant *Rumex hastatulus*. *Proceedings of the National Academy of Sciences of the United States of America* **111**(21): 7713–7718.
- Hughes JF, Skaletsky H, Pyntikova T, Graves TA, Daalen SKMv, Minx PJ, Fulton RS, McGrath SD, Locke DP, C.Friedman, Trask BJ, Mardis ER, Warren WC, Repping S, Rozen S, Wilson RK, Page DC. 2010.** Chimpanzee and human Y chromosomes are remarkably divergent in structure and gene content. *Nature* **463**: 536-539.
- Immler S, Otto S. 2015.** The evolution of sex chromosomes in organisms with separate haploid sexes. *Evolution* **69**(3): 694–708.
- Iovene M, Yu Q, Ming R, Jiang J. 2015.** Evidence for emergence of sex-determining gene(s) in a centromeric region in *Vasconcellea parviflora*. *Genetics* **199**: 413-421.

- 896 **Jakse J, Stajner N, Kozjak P, Cerenak A, Javornik B. 2008.** Trinucleotide
897 microsatellite repeat is tightly linked to male sex in hop (*Humulus lupulus* L.).
898 *Molecular Breeding* **21**: 139-148.
- 899 **Jordan C, Charlesworth D. 2012.** The potential for sexually antagonistic
900 polymorphism in different genome regions. *Evolution* **66**(2): 505–516.
- 901 **Khattak J, Torp A, Andersen S. 2006.** A genetic linkage map of *Spinacia oleracea*
902 and localization of a sex determination locus. *Euphytica* **148**(3): 311-318.
- 903 **Kubat Z, Zluvova J, Vogel I, Kovacova V, Cermak T, Cegan R, Hobza R, Vyskot
904 B, Kejnovsky E. 2014.** Possible mechanisms responsible for absence of a
905 retrotransposon family on a plant Y chromosome. *New Phytologist* **202**: 662-
906 678.
- 907 **Lahn BT, Page DC. 1999.** Four evolutionary strata on the human X chromosome.
908 *Science* **286**: 964-967.
- 909 **Lardon A, Georgiev S, Aghmir A, Merrer GL, Negrutiu I. 1999.** Sexual
910 dimorphism in white campion: complex control of carpel number is revealed
911 by Y chromosome deletions. *Genetics* **151**: 1173-1185.
- 912 **Lindholm A, Breden F. 2002.** Sex chromosomes and sexual selection in Poeciliid
913 fishes. *Am. Nat.* **160**: S214-S224.
- 914 **Liu Z, Moore PH, Ma H, Ackerman CM, Ragiba M, Pearl HM, Kim MS,
915 Charlton JW, Yu Q, Stiles JI, Zee FT, Paterson AH, Ming R. 2004.** A
916 primitive Y chromosome in Papaya marks the beginning of sex chromosome
917 evolution. *Nature* **427**: 348-352.
- 918 **Lloyd DG. 1982.** Selection of combined versus separate sexes in seed plants. *Amer.*
919 *Nat.* **120**: 571-585.
- 920 **Maddison WP, Leduc-Robert G. 2013.** Multiple origins of sex chromosome fusions
921 correlated with chiasma localization in Habronattus jumping spiders (Araneae:
922 Salticidae) *Evolution* **67**(8): 2258–2272.
- 923 **Mamanova L, Coffey A, Scott CE, Kozarewa I, Turner E, Kumar A, Howard E,
924 Shendure J, Turner D. 2010.** Target-enrichment strategies for next-
925 generation sequencing. *Nature Methods* **7**(2): 111-118.
- 926 **Marais GAB, Forrest A, Kamau E, Käfer J, Daubin V, Charlesworth D. 2011.**
927 Multiple nuclear gene phylogenetic analysis of the evolution of dioecy and sex
928 chromosomes in the genus *Silene*. *PLOS ONE* **6**(8): e21915.
- 929 **Mariotti B, Manzano S, Kejnovsky E, Vyskot B, Jamilena M. 2008.** Accumulation
930 of Y-specific satellite DNAs during the evolution of *Rumex acetosa* sex
931 chromosomes. *Molecular Genetics and Genomics* **281**: 249-259.
- 932 **Matsunaga S, Isono E, Kejnovsky E, Vyskot B, Kawano S, Charlesworth D.
933 2003.** Duplicative transfer of a MADS box gene to a plant Y chromosome.
934 *Molecular Biology and Evolution* **20**(7): 1062-1069.
- 935 **McDaniel SF, Neubig KM, Payton AC, Quatrano RS, Cove DJ. 2013.** Recent
936 gene-capture on the UV sex chromosomes of the moss *Ceratodon purpureus*.
937 *Evolution* **67**(10): 2811–2822.
- 938 **McDaniel SF, Willis JH, Shaw AJ. 2007.** A linkage map reveals a complex basis for
939 segregation distortion in an interpopulation cross in the moss *Ceratodon*
940 *purpureus*. *Genetics* **176**: 2489-2500.
- 941 **Ming R, Bendahmane A, Renner S. 2011.** Sex chromosomes in land plants.
942 *ANNUAL REVIEW OF PLant Biology* **62**: 485-514.
- 943 **Mitchell C, Diggle P. 2005.** The evolution of unisexual flowers: morphological and
944 functional convergence results from diverse developmental transitions.
945 *American Journal of Botany* **92**: 1068-1076.

- Mrackova M, Nicolas M, Hobza R, Negrutiu I, Monéger F, Widmer A, Vyskot B, Janousek B. 2008. Independent origin of sex chromosomes in two species of the genus *Silene*. *Genetics* **179**: 1129-1133.
- Muller HJ. 1932. Some genetic aspects of sex. *The American Naturalist* **66**: 118-138.
- Muller HJ. 1950. Evidence of the precision of genetic adaptation. *Harvey Lectures* **43**: 165-229.
- Muyle A, Zemp N, Deschamps C, Mousset S, Widmer A, Marais G. 2012. Rapid de novo evolution of X chromosome dosage compensation in *Silene latifolia*, a plant with young sex chromosomes. *PLoS Biology* **10**(4): e1001308.
- Myosho T, H Otake HM, Masaru Matsuda, Yoko Kuroki, Asao Fujiyama, Kiyoshi Naruse, Satoshi Hamaguchi, Sakaizumi M. 2012. Tracing the emergence of a novel sex-determining gene in medaka, *Oryzias luzonensis*. *Genetics* **191**: 163-170.
- Navajas-Pérez R, Herran Rdl, Gonzalez G, Jamilena M, Lozano R, Rejon C, Rejon M, Garrido-Ramos MA. 2005. The evolution of reproductive systems and sex-determining mechanisms within *Rumex* (Polygonaceae) inferred from nuclear and chloroplastidial sequence data. *Molecular Biology and Evolution* **22**(9): 1929-1193.
- Nicolas M, Marais G, Hykelova V, Janousek B, Laporte V, Vyskot B, Mouchiroud D, Negrutiu I, Charlesworth D, Monéger F. 2005. A gradual process of recombination restriction in the evolutionary history of the sex chromosomes in dioecious plants. *PLoS Biology* **3**(1): 47-56.
- Nishiyama R, Ishii K, Kifune E, Kazama Y, Nishihara K, S M, Shinozaki K, Kawano S. 2010. Sex chromosome evolution revealed by physical mapping of *SIAP3X/Y* in the dioecious plant *Silene latifolia* *Cytologia* **75**(3): 319-325.
- Ohnishi O. 1985. Population genetics of cultivated buckwheat, *Fagopyrum esculentum* Moench. III. Frequency of sterility mutants in Japanese populations. *Japanese Journal of Genetics* **60**: 391-404.
- Okada S, Sone T, Fujisawa M, Nakayama S, Takenaka M, Ishizaki K, Kono K, Shimizu-Ueda Y, Hanajiri T, Yamato KT, Fukuzawa H, Brennicke A, Ohyama K. 2001. The Y chromosome in the liverwort *Marchantia polymorpha* has accumulated unique repeat sequences harboring a male-specific gene. *Proceedings of the National Academy of Sciences of the United States of America* **98**: 9454-9459.
- Onodera Y, Itaru Y, Hiroki M, Tanaka A, Niikura S, Yamazaki S, Mikami T. 2011. Mapping of the genes for dioecism and monoecism in *Spinacia oleracea* L.: evidence that both genes are closely linked. *Plant Cell Reports* **30**(6): 965-971.
- Oyama R, Volz S, Renner S. 2009. A sex-linked SCAR marker in *Bryonia dioica* (Cucurbitaceae), a dioecious species with XY sex-determination and homomorphic sex chromosomes. *Journal of Evolutionary Biology* **22**(1): 214-224.
- Pakull B, Groppe K, Mecucci F, Gaudet M, Sabatti M, Fladung M. 2011. Genetic mapping of linkage group XIX and identification of sex-linked SSR markers in a *Populus tremula* × *Populus tremuloides* cross. *Canadian Journal of Forest Research* **41**(2): 245-253.
- Pane A, Simone AD, Saccone G, Polito C. 2005. Evolutionary conservation of *Ceratitis capitata* transformer gene function. *Genetics* **171**(2): 615-624.
- Pannell J. 1997. Mixed genetic and environmental sex determination in an androdioecious population of *Mercurialis annua*. *Heredity* **78**: 50-56.

- 996 **Park J, Ishikawa Y, Yoshida R, Kanno A, Kameya T. 2003.** Expression of
997 AODEF, a B functional MADS-box gene, in stamens and inner tepals of the
998 dioecious species *Asparagus officinalis* L. *Plant Molecular Biology* **51**(6):
999 867-875.
- 1000 **Peil A, Flachowsky H, Schumann E, Weber WE. 2003.** Sex-linked AFLP markers
1001 indicate a pseudoautosomal region in hemp (*Cannabis sativa* L.). *Theoretical*
1002 *and Applied Genetics* **107**(1): 102-109.
- 1003 **Penny R, Montgomery B, Delph L. 2011.** X Linkage of AP3A, a Homolog of the Y-
1004 Linked MADS-Box Gene *AP3Y* in *Silene latifolia* and *S. dioica*. *PLOS ONE*
1005 **6**(4): e18972.
- 1006 **Picq S, Santoni S, Lacombe T, Latreille M, Weber A, Ardisson M, Ivorra S,**
1007 **Maghradze D, Garcia-Arroyo R, This P, Terral J-F, Bacilieri R, A small**
1008 **XY chromosomal region explains sex determination in wild dioecious *V.***
1009 ***vinifera* and the reversal to hermaphroditism in domesticated grapevines.**
1010 **BMC plant biology. 2014.** A small XY chromosomal region explains sex
1011 determination in wild dioecious *V. vinifera* and the reversal to
1012 hermaphroditism in domesticated grapevines. *BMC Plant Biology* **14**: 229.
- 1013 **Pigozzi M. 2011.** Diverse stages of sex-chromosome differentiation in tinamid birds:
1014 evidence from crossover analysis in *Eudromia elegans* and *Crypturellus*
1015 *tataupa*. *Genetica* **139**(6): 771-777.
- 1016 **Policansky D. 1981.** Sex choice and the size advantage model in jack-in-the-pulpit.
1017 *Proc. Natl. Acad. Sci. USA* **78**: 1306-1308.
- 1018 **Pomiankowski A, Nothiger R, Wilkins A. 2004.** The evolution of the *Drosophila*
1019 sex-determination pathway. *Genetics* **166**: 1761-1773.
- 1020 **Ramos M, Coito J, Silva H, Cunha J, Costa M, Rocheta M. 2014.** Flower
1021 development and sex specification in wild grapevine *BMC Genomics* **15**:
1022 1095.
- 1023 **Renner S. 2014.** The relative and absolute frequencies of angiosperm sexual systems:
1024 Dioecy, monoecy, gynodioecy, and an updated online database. *American*
1025 *Journal of Botany* **101**: 1588-1596.
- 1026 **Renner SS, Ricklefs RE. 1995.** Dioecy and its correlates in the flowering plants.
1027 *American Journal of Botany* **82**: 596-606.
- 1028 **Renner SS, Won H. 2001.** Repeated evolution of dioecy from monoecy in
1029 Siparunaceae (Laurales). *Systematic Biology* **50**(5): 700-712.
- 1030 **Rice WR. 1987.** The accumulation of sexually antagonistic genes as a selective agent
1031 promoting the evolution of reduced recombination between primitive sex-
1032 chromosomes. *Evolution* **41**: 911-914.
- 1033 **Rodgers-Melnicka E, Bradbury PJ, Elshire RJ, Glaubitz JC, Acharya CB,**
1034 **Mitchella SE, Li C, Lic Y, Buckler ES. 2015.** Recombination in diverse
1035 maize is stable, predictable, and associated with genetic load. *Proceedings of*
1036 *the National Academy of Sciences of the United States of America* **112**(5):
1037 3823-3828.
- 1038 **Ross JA, Peichel CL. 2008.** Molecular cytogenetic evidence of rearrangements on
1039 the Y chromosome of the threespine stickleback fish. *Genetics* **179**(4): 2173-
1040 2182.
- 1041 **Ross JA, Urton JR, Boland J, Shapiro MD, Peichel CL. 2009.** Turnover of sex
1042 chromosomes in the stickleback fishes (Gasterosteidae). *PLoS Genetics* **5**:
1043 e1000391.
- 1044 **Saccone G, Pane A, Polito LC. 2002.** Sex determination in flies, fruitflies and
1045 butterflies. *Genetica* **116**(1): 15-23.

- 1046 **Sakamoto K, Abe T, Matsuyama T, Yoshida S, Ohmido N, Fukui K, Satoh S.**
 1047 **2005.** RAPID markers encoding retrotransposable elements are linked to the
 1048 male sex in *Cannabis sativa* L. *Genome* **48**(5): 931-936.
- 1049 **Scotti I, Delph LF. 2006.** Selective trade-offs and sex-chromosome evolution in
 1050 *Silene latifolia*. *Evolution* **60**(9): 1793–1800.
- 1051 **Shibata F, Hizume M, Kuroki Y. 2000.** Differentiation and the polymorphic nature
 1052 of the Y chromosomes revealed by repetitive sequences in the dioecious plant,
 1053 *Rumex acetosa*. *Chromosome Research* **8**: 229-236.
- 1054 **Siljak-Yakovlev S, Benmalek S, Cerbah M, Peña TCdl, Bounaga N, Brown SC,**
 1055 **Sarr A. 1996.** Chromosomal sex determination and heterochromatin structure
 1056 in date palm. *Sexual Plant Reproduction* **9**: 127-132.
- 1057 **Skaletsky H, Kuroda-Kawaguchi T, Minx PJ, Cordum HS, Hillier L, Brown LG,**
 1058 **Repping S, Pyntikova T, Ali J, Bieri T, Chinwalla A, Delehaunty A,**
 1059 **Delehaunty K, Du H, Fewell G, Fulton L, Fulton R, Graves T, Hou S,**
 1060 **Latrielle P, Leonard S, Mardis E, Maupin R, McPherson J, Miner T,**
 1061 **Nash W, Nguyen C, Ozersky P, Pepin K, Rock S, Rohlfsing T, Scott K,**
 1062 **Schultz B, Strong C, Tin-Wollam A, Yang S, Waterston RH, Wilson RK,**
 1063 **Rozen S, Page DC. 2003.** The male-specific region of the human Y
 1064 chromosome is a mosaic of discrete sequence classes. *Nature* **423**: 825 - 837.
- 1065 **Smith BW. 1964.** The evolving karyotype of *Rumex hastatulus*. *Evolution* **18**: 93-
 1066 104.
- 1067 **Sousa A, Fuchs J, Renner SS. 2012.** Molecular cytogenetics (FISH, GISH) of
 1068 *Coccinia grandis*: A ca. 3 myr-old species of Cucurbitaceae with the Largest
 1069 Y/autosome divergence in flowering plants. *Cytogenetic and Genome*
 1070 *Research* **139**(2): 107-118.
- 1071 **Spigler R, Lewers K, Main D, Ashman T-L. 2008.** Genetic mapping of sex
 1072 determination in a wild strawberry, *Fragaria virginiana*, reveals earliest form
 1073 of sex chromosome. *Heredity* **101**(6): 507-517.
- 1074 **Suetsugu Y, Futahashi R, Kanamori H, Kadono-Okuda K, Sasanuma S-i,**
 1075 **Narukawa J, Ajimura M, Jouraku A, Namiki N, Shimomura M, Sezutsu**
 1076 **H, Osanai-Futahashi M, Suzuki MG, Daimon T, Shinoda T, Taniai K,**
 1077 **Asaoka K, R Niwa, Kawaoka S, Katsuma S, Tamura T, Noda H,**
 1078 **Kasahara M, Sugano S, Y Suzuki, Fujiwara H, Kataoka H, Arunkuma**
 1079 **KP, Tomar A, Nagaraju J, Goldsmith MR, Feng Q, Xia Q, Yamamoto K,**
 1080 **Shimada T, Mita K. 2013.** Large scale full-length cDNA sequencing reveals
 1081 a unique genomic landscape in a Lepidopteran model insect, *Bombyx mori*. *G3*
 1082 **3**: 1481-1492.
- 1083 **Tanksley SD, Zamir D, Rick CM. 1981.** Evidence for extensive overlap of
 1084 sporophytic and gametophytic gene expression in *Lycopersicon esculentum*.
 1085 *Science* **213**: 453-455.
- 1086 **Telgmann-Rauber A, Jamsari A, Kinney MS, Pires JC, Jung C. 2007.** Genetic
 1087 and physical maps around the sex-determining M-locus of the dioecious plant
 1088 asparagus. *Molecular Genetics and Genomics* **278**: 221-234.
- 1089 **Tennessen J, Govindarajul R, Liston A, Ashman T. 2013.** Targeted sequence
 1090 capture provides insight into genome structure and genetics of male sterility in
 1091 a gynodioecious diploid strawberry, *Fragaria vesca* ssp. *bracteata* (Rosaceae).
 1092 *G3* **3**: 1341-1351.
- 1093 **Ubisch GV. 1936.** Genetic studies on the nature of hermaphroditic plants in
 1094 *Antennaria dioica* (L) Gaertn. *Genetics* **21**(3): 282-294.

- 1095 **Uno Y, Nishida C, Oshima Y, Yokoyama S, Miura I, Matsuda Y, Nakamura M.**
 1096 **2008.** Comparative chromosome mapping of sex-linked genes and
 1097 identification of sex chromosomal rearrangements in the Japanese wrinkled
 1098 frog (*Rana rugosa*, Ranidae) with ZW and XY sex chromosome systems.
 1099 *Chromosome Research* **16**(4): 637-647.
- 1100 **Van Buren R, Zeng F, Chen C, Zhang J, Wai C, Han J, Aryal R, Gschwend AR,**
 1101 **Wang J, Na J, Huang L, Zhang L, Miao W, Gou J, Arro J, Guyot R,**
 1102 **Moore RC, Wang M, Zee F, Charlesworth D, Moore PH, Yu Q, Ming R.**
 1103 **2015.** Origin and domestication of papaya Yh chromosome. *Genome*
 1104 *Research*.
- 1105 **van Doorn G, Kirkpatrick M. 2007.** Turnover of sex chromosomes induced by
 1106 sexual conflict. *Nature* **449**: 909-912.
- 1107 **Vicoso B, Charlesworth B. 2006.** Evolution on the X chromosome: unusual patterns
 1108 and processes. *Nature Reviews Genetics* **7**: 645-653.
- 1109 **Vicoso B, Kaiser V, Bachtrog D. 2013.** Sex-biased gene expression at homomorphic
 1110 sex chromosomes in emus and its implication for sex chromosome evolution.
 1111 *Proceedings of the National Academy of Sciences of the United States of*
 1112 *America* **110**(16): 6453-6458.
- 1113 **Vuilleumier S, Lande R, JJM VA, Seehausen O. 2007.** Invasion and fixation of
 1114 sex-reversal genes *Journal of Evolutionary Biology* **20**(3): 913-920.
- 1115 **Wang J, Na J, Yu Q, Gschwend AR, Han J, Zeng F, Aryal R, Van Buren R,**
 1116 **Murray JE, Zhang W, Navajas-Pérez R, Feltus F, Lemke C, Tong EJ,**
 1117 **Chen C, Wai C, Singh R, Wang M, Min X, Alam M, Charlesworth D,**
 1118 **Moore PH, Jiang J, Paterson AH, Ming R. 2012.** Sequencing papaya X and
 1119 Yh chromosomes reveals molecular basis of incipient sex chromosome
 1120 evolution. *Proceedings of the National Academy of Sciences of the United*
 1121 *States of America* **109**(34): 13710-13715.
- 1122 **Webb C. 1979.** Breeding system and seed set in *Euonymus europaeus* (Celastraceae).
 1123 *Plant Systematics and Evolution* **132**(4): 299-303.
- 1124 **Wellmer F, Alves-Ferreira M, Dubois A, Riechmann J, Meyerowitz E. 2006.**
 1125 Genome-wide analysis of gene expression during early Arabidopsis flower
 1126 development. *PLoS Genetics* **2**(7): e117.
- 1127 **Westergaard M. 1958.** The mechanism of sex determination in dioecious plants.
 1128 *Advances in Genetics* **9**: 217-281.
- 1129 **Wilkins A. 1995.** Moving up the hierarchy: a hypothesis on the evolution of a genetic
 1130 sex-determination pathway. *Bioessays* **17**(1): 71-77.
- 1131 **Yamato KT, Ishizaki K, Fujisawa M, Okada S, Nakayama S, Fujishita M, Bando**
 1132 **H, Yodoya K, Hayashi K, Bando T, Hasumi A, Nishio T, Sakata R,**
 1133 **Yamamoto M, Yamaki M, Kajikawa M, Yamano T, Nishide T, Choi SH,**
 1134 **Shimizu-Ueda Y, Hanajiri T, Sakaida M, Kono K, Takenaka M,**
 1135 **Yamaoka S, Kuriyama C, Kohzu Y, Nishida H, Brennicke A, Shin-I T,**
 1136 **Kohara Y, Kohchi T, Fukuzawa H, Ohyama K. 2007.** Gene organization of
 1137 the liverwort Y chromosome reveals distinct sex chromosome evolution in a
 1138 haploid system. *Proceedings of the National Academy of Sciences of the USA*
 1139 **104**: 6472-6477.
- 1140 **Yin T, DiFazio S, Gunter L, Zhang X, Sewell M, Woolbright S, Allan G, Kelleher**
 1141 **C, Douglas C, Wang M, Tuskan G. 2008.** Genome structure and emerging
 1142 evidence of an incipient sex chromosome in *Populus*. *Genome Research* **18**:
 1143 422-430.

- 1144 **Yoshida K, T Makino T YK, Shigenobu S, Peiche C, Toyoda A, Fujiyama A,**
 1145 **Kitano J. 2014.** Sex Chromosome Turnover Contributes to Genomic
 1146 Divergence between Incipient Stickleback Species. *PLoS Genetics* **10**(3):
 1147 e1004223.
- 1148 **Yu Q, Hou S, Hobza R, Feltus F, Wang X, Jin W, Skelton R, Blas A, Lemke C,**
 1149 **Saw J, Moore P, Alam M, Jiang J, Paterson A, Vyskot B, Ming R. 2007.**
 1150 Chromosomal location and gene paucity of the male specific region on papaya
 1151 Y chromosome. *Molecular Genetics and Genomics* **278**(2): 177-185.
- 1152 **Zhang W, Wang X, Yu Q, Ming R, Jiang J. 2008.** DNA methylation and
 1153 heterochromatinization in the male-specific region of the primitive Y
 1154 chromosome of papaya. *Genome Research*(18).
- 1155 **Zhou Q, Bachtrog D. 2012.** Chromosome-wide gene silencing initiates Y
 1156 degeneration in *Drosophila*. *Current Biology* **22**: 522–525.
- 1157 **Zhou Q, Zhang J, Bachtrog D, An N, Huang Q, Jarvis ED, Gilbert MTP, Zhang**
 1158 **G. 2014.** Complex evolutionary trajectories of sex chromosomes across bird
 1159 taxa. *Science* **346**: 1332.
- 1160 **Zimmerman JK. 1991.** Ecological correlates of labile sex expression in the orchid
 1161 *Catasetum viridiflavum* *Ecology* **72**(2): 597-608.
- 1162 **Zluvova J, Janousek B, Negrutiu I, Vyskot B. 2005.** Comparison of the X and Y
 1163 chromosome organization in *Silene latifolia*. *Genetics* **170**: 1431-1434.
- 1164
 1165
 1166

1167 **Figure legends**

1168

1169

1170 **Figure 1.** Possible histories of sex-linked regions. A. Sex-determining genome region
1171 with no prior history of suppressed recombination, in which recombination
1172 suppression evolves in response to a polymorphism at two sex-determining
1173 loci. B. Pericentromeric region with suppressed recombination, in which
1174 sex-determining loci evolve. The sex-linked region may later become
1175 extended by evolution of a further region of suppressed recombination,
1176 perhaps due to the establishment of a sexually antagonistic polymorphism
1177 on the same chromosome (see text). C. A take-over event in which a new
1178 male-determining gene arises in a genome region not previously involved
1179 in sex-determination; the figure illustrates the case when the new sex-
1180 determining gene is on a chromosome (indicated by a dashed line) different
1181 from the previously existing one, but a changed location on the same
1182 chromosome is also possible.

1183

1184 **Figure 2.** Evolution of sex-determining and sex-linked genome regions. A. Evolution
1185 of sex-determining genes in a genome region starting from an
1186 hermaphrodite ancestor, and of suppressed recombination in the region,
1187 forming a sex chromosome-like region, showing disadvantageous
1188 recombinants between the proto-Y and the proto-X chromosomes (the
1189 reciprocal recombinant would be hermaphroditic, and is not shown). If the
1190 female suppressor has male-specific expression (or evolves expression
1191 restricted to males), it can spread through the entire population, and create a
1192 single-gene sex-determining system (bottom left). B. Evolution of sex-
1193 determining genes in a monoecious ancestor.

1194

1195 **Figure 3.** Loss of genes from sex-linked regions. A. Chromosome before gene loss.
1196 The region that will evolve sex linkage includes four genes essential for
1197 vegetative functions in both sexes (green), and other genes essential only in
1198 males (blue) or females (pink), or affecting both sex functions (both
1199 colours). B. Loss of female function genes from a Y chromosome in a
1200 diploid plant, and loss of the third gene essential vegetative functions


1201 (which is non-lethal due to the presence of the copy on the X chromosome).
1202 C. Loss of female and male function genes from male- and female-
1203 determining chromosomes in a haploid plant. Some genes with functions in
1204 the diploid vegetative stage could also be lost from either the U or the V.
1205

For Peer Review

Table 1. Some characteristics favourable for studying the genetics and evolution of plant sex chromosomes. The main text provides examples from plant studies.

Characteristic	Advantages	Specific evolutionary questions
A range of sex chromosome ages exist, including recently evolved ones and old-established systems	The time when recombination stopped can be estimated using a molecular clock, as it is often not long, and sequence differences will not be saturated, but will reflect times when recombination was suppressed	1. Which species without cytologically visible heteromorphism have sex-linked regions that include genes other than the sex-determining genes?
		2. Does recombination suppression always evolve, even in old-established systems, or does it sometimes fail to evolve (and, if so, why)?
	The earliest stages of sex chromosome evolution can be studied	1. Is there a tendency for chromosomal heteromorphism, heterochromatinisation, ZW systems, and X-autosome balance systems to be associated with older-established systems?
		2. How did recombination suppression evolve (gradually, or in distinct recombination suppression events affecting genome regions with many genes), and how often do such events happen?
Closely related non-dioecious outgroup species often exist	The directions of changes during sex chromosome evolution can be studied	3. Did repetitive sequences accumulate before genes started to lose functions, or does their accumulation contribute to loss of functions?
		1. Have plant X and/or Y chromosomes adapted to the new dioecious state?
Dioecy evolved repeatedly	The directions of changes during sex chromosome evolution can be studied	2. Have plant Y chromosomes degenerated genetically? If so, what is the time course?
		In taxa with many dioecious species, is dioecy ancestral or has it evolved several times; if the latter, are the same sex-determining genes involved, or different genes in different species?

Figure 1



KEY



Centromere



Sex determining genes

Regions with suppressed recombination



Pericentromeric region



Male-specific region

Figure 2




Figure 2, continued

B Initially monoecious cosexual





Figure 3

A



B



C

